

Algebraic Ranks of CAT(0) Groups*

Raeyong Kim

October 8, 2012

Abstract

We study the algebraic rank of various classes of CAT(0) groups. They include right-angled Coxeter groups, right-angled Artin groups, relatively hyperbolic groups and groups acting geometrically on CAT(0) spaces with isolated flats. As one of our corollaries, we obtain a new proof of a result on commensurability of Coxeter groups.

1 Introduction

Let M be a complete Riemannian manifold of nonpositive sectional curvature. The *geometric rank* of a geodesic γ in M , denoted by $rk(\gamma)$, is the dimension of the vector space of parallel Jacobi fields along γ . Then the *geometric rank* of M is defined to be the minimum of $rk(\gamma)$ over all geodesics γ in M .

The celebrated rank-rigidity theorem, due to Ballmann ([1]), and to Burns-Spatzier ([4]), states that if M has bounded nonpositive sectional curvature and finite volume, then the universal cover \widetilde{M} is a flat Euclidean space, a symmetric space of non-compact type, a space of rank 1 or a product of such spaces.

In [26], Prasad and Raghunathan introduced the notion of the algebraic rank, $\text{rank}(G)$, of a group G . (See Section 2 for the definition.) Ballmann and Eberlein proved that if Γ is the fundamental group of a complete Riemannian manifold M of bounded nonpositive sectional curvature and of finite volume, then $\text{rank}(\Gamma)$ is equal to the geometric rank of M (see [2]). By combining this with the rank-rigidity theorem, we have that, for a complete Riemannian manifold M of bounded nonpositive sectional curvature and of finite volume, if \widetilde{M} does not have an Euclidean factor and $\Gamma = \pi_1(M)$ has higher algebraic rank, then either Γ has a finite index subgroup which splits as a direct product or it is a lattice in a semi-simple Lie group of higher rank.

There is an analogous notion of geometric rank for CAT(0) spaces. A *geometric flat of dimension n* in a complete CAT(0) space X is a closed convex subset of X which is isometric to the Euclidean n -space. A geodesic line L is said to have *rank one* if it does not bound a flat half-plane. A complete CAT(0) space X is said to have *higher geometric rank* if no geodesic in X has rank one.

*2010 *AMS Subject Classification* 57M07 (20F55, 20F67, 20F65)

Let X be a complete CAT(0) space and G be a group acting geometrically (i.e., properly and cocompactly by isometries) on X . In view of the Ballmann-Eberlein result, it is natural to ask the similar question for CAT(0) spaces:

Conjecture 1 G has higher algebraic rank if and only if X has higher geometric rank.

In this paper, we study the algebraic rank of various CAT(0) groups. They include right-angled Coxeter groups, right-angled Artin groups, relatively hyperbolic groups and groups acting geometrically on CAT(0) spaces with isolated flats. In Section 3, we prove that if W is an infinite irreducible non-affine right-angled Coxeter group, then $\text{rank}(W) = 1$. As a corollary, we obtain a new proof for the question posed by M. Davis in [7], namely, W cannot be commensurable to any uniform lattice in a higher rank non-compact connected semi-simple Lie group. In Subsection 3.2, we prove that any non-join right-angled Artin group has an algebraic rank of 1. In Section 4, we study algebraic ranks of groups which act on CAT(0) spaces with isolated flats. More precisely, if a group G acts geometrically on CAT(0) space with isolated flats \mathcal{F} and $|\mathcal{F}| \neq 1$, then $\text{rank}(G) \leq 1$. It is well known that such a group G is hyperbolic relative to a family of stabilizers of flats in \mathcal{F} . We study the dynamics of a relatively hyperbolic group acting on the boundary of a δ -hyperbolic space to prove that the algebraic rank of a relative hyperbolic group is ≤ 1 if there are at least two peripheral subgroups containing elements of infinite order. It follows immediately that $\text{rank}(G) \leq 1$.

This paper is part of author's Ph.D. thesis. The author thanks the thesis advisor Jean Lafont for his guidance throughout this research project. The author also thanks Mike Davis for helpful conversations concerning Coxeter groups.

2 Algebraic Ranks of Groups

Definition 2 For a given group G , let $\mathcal{A}_i(G)$ be the set consisting of elements such that the centralizer contains a free abelian subgroup of rank $\leq i$ as a subgroup of finite index. Define $r(G)$ to be the minimum i such that G can be expressed as the union of finitely many translates of $\mathcal{A}_i(G)$. In other words,

$$r(G) = \min\{i \mid \text{there are finitely many elements } g_j \in G \text{ such that } G = \bigcup_j g_j \mathcal{A}_i(G)\}$$

Finally, the *algebraic rank* of G , $\text{rank}(G)$, is the supremum of $r(G^*)$ over all finite index subgroups G^* of G .

We allow the possibility that $\text{rank}(G) = 0$. For example, if G is a finite group, then $\text{rank}(G) = 0$. On the other hand, if G is torsion-free, then $\text{rank}(G) > 0$: suppose that $\text{rank}(G) = 0$. In particular, $r(G) = 0$. Then the set $\mathcal{A}_0(G)$ must be non-empty. But $\mathcal{A}_0(G)$ is a subset of the set of finite order elements in G .

We set $\text{rank}(G) = \infty$ if the sets $\mathcal{A}_i(G)$ are empty, or if G can not be covered by finitely many translates of any of the sets $\mathcal{A}_i(G)$. For example, if a group G has an infinitely generated free abelian center, then $\text{rank}(G) = \infty$. In fact, there

exist finitely presented examples of such groups. More specifically, Hall obtained in [16] the existence of a finitely generated group which has infinitely generated free abelian center. Using [25], we can obtain a finitely presented group having infinitely generated free abelian center.

Remark 3 $r(G)$ is not necessarily equal to $\text{rank}(G)$. Following [2], we present an example of a group satisfying $r(G) < \text{rank}(G)$. See [2, Section 4] for more examples.

Let G be the fundamental group of a flat Klein bottle, acting on \mathbb{E}^2 by isometries. (In the simplest case G is generated by $\phi_1 : (x, y) \rightarrow (x + 1, -y)$ and $\phi_2 : (x, y) \rightarrow (x, y + 1)$.) The set $\mathcal{A}_1(G)$ consists of all elements of G that reverse the orientation of \mathbb{E}^2 . Then $G = \mathcal{A}_1(G) \cup \gamma \mathcal{A}_1(G)$, for any $\gamma \in \mathcal{A}_1(G)$. Therefore, $r(G) < 2 = \text{rank}(G) = \text{rank}(\mathbb{E}^2)$.

We close the section by mentioning that algebraic ranks of groups behave well under products and taking finite index subgroups.

Proposition 4 [2, Proposition 2.1] *Let G be an abstract group.*

1. *If G' is a finite index subgroup of G , then $r(G) \leq r(G')$ and $\text{rank}(G) = \text{rank}(G')$.*
2. *If $G = G_1 \times \cdots \times G_n$, then*

$$r(G) = \sum_{i=1}^n r(G_i), \quad \text{rank}(G) = \sum_{i=1}^n \text{rank}(G_i)$$

3 Right -Angled Coxeter Groups and Artin Groups

3.1 Coxeter Groups

A *Coxeter system* (W, S) is a group W and a set $S = \{s_1, s_2, \dots\}$ of generators such that W has the following presentation

$$W = \langle S \mid (s_i s_j)^{m_{ij}} = 1, s_i, s_j \in S \rangle,$$

where $m_{ii} = 1$ and if $i \neq j$, then $m_{ij} = m_{ji}$ is a positive integer ≥ 2 or ∞ (in which case we omit the relation between s_i and s_j). W is called a *Coxeter group*. The cardinality $|S|$ of S is called the *rank* of W and we assume that $|S|$ is finite in this section. A Coxeter group W is *spherical* if W is finite and *affine* if W has a finite index free abelian subgroup.

For any subset $J \subset S$, we denote by W_J the subgroup of W generated by J . We call W_J a *standard parabolic subgroup*, and any conjugate of a standard parabolic subgroup is called a *parabolic subgroup*. For any subset $A \subset W$, the *parabolic closure* $\text{Pc}(A)$ of A is the smallest parabolic subgroup containing A . An element γ is called *essential* if $\text{Pc}(\gamma) = W$.

Associated to any Coxeter group W , there is a CAT(0) polyhedral cell complex Σ_W , which is called the *Davis complex*, upon which W acts properly discontinuously and cocompactly by isometries. Σ_W can be cellulated by so called *Coxeter polytopes* and, with a natural Euclidean metric on each Coxeter polytope, inherits a piecewise Euclidean metric. It was Gromov (right-angled case, [15]) and Mousong (general case, [23]), who showed that Σ_W , with this metric, is CAT(0). See [8] for details. An element $\gamma \in W$ is said to have *rank one* if it is hyperbolic and if some (and hence any) of its axes in Σ_W has rank one. In [5], Caprace and Fujiwara study rank one elements in Coxeter groups. In particular, an element γ has rank one if and only if its centralizer is virtually infinite cyclic. In other words, any rank one element is contained in $\mathcal{A}_1(W)$.

Right-angled Coxeter groups are Coxeter groups for which $m_{ij} = 2$ or ∞ for $i \neq j$. In this case, the Davis complex Σ_W is a CAT(0) cubical complex. In this subsection, we prove that any infinite irreducible non-affine right-angled Coxeter group has an algebraic rank of 1.

Remark 5 1. Any spherical Coxeter group has an algebraic rank of 0. (See Section 2.)

2. It is a consequence of Selberg's lemma that every infinite Coxeter group W has a torsion-free subgroup of finite index. Therefore, such groups satisfy $\text{rank}(W) \geq 1$.
3. If W is infinite, irreducible and affine, then $\text{rank}(W) = |S| - 1$.
4. Suppose that W is infinite and reducible, $W = W_{T_1} \times W_{T_2} \times \cdots \times W_{T_n}$. By Proposition 4, $\text{rank}(W) = \sum_{i=1}^n \text{rank}(W_{T_i})$. Therefore, $\text{rank}(W) > 1$ unless W is virtually cyclic.

Hereafter, we assume that W is an infinite irreducible non-affine right-angled Coxeter group. Tits' solution to the word problem for Coxeter groups states that any two reduced expressions represent the same element in W if and only if one can be transformed into the other by a series of replacements of the alternating subword $(stst\cdots)$ of length m_{st} by the subword $(tsts\cdots)$ of length m_{st} . (See [8, Sec. 3.4].) This implies

Lemma 6 *For $w \in W$, let $S(w)$ be the set of generators appearing in some (and hence any) reduced expression for w . If $s \in S(w)$ appears an odd (respectively, even) number of times in some reduced expression for w , then s appears an odd (respectively, even) number of times in any reduced expression for w .*

Let \mathcal{H} be the set of rank one elements in W . It is not difficult to find rank one elements in W . For example, any essential element in W is virtually infinite cyclic, therefore it has rank one. (See [21, Corollary 6.3.10])

Lemma 7 *Let w be an element such that some (and hence any) reduced expression for w has the following property : all generators appear, and each generator appears an odd number of times. Then w is essential, and hence, w has rank one.*

Proof. Suppose that $\text{Pc}(w) = uW_Ju^{-1}$ for some u and some $J \subset S$. Suppose that $s \notin J$. Then any reduced expression for words in uW_Ju^{-1} contains s an even number of times. By Lemma 6, $s \in J$. This proves that $J = S$. ■

Proposition 8 $r(W) \leq 1$.

Proof. Recall $\mathcal{H} \subset \mathcal{A}_1(W)$. Let $\mathcal{S} = \{s_1 \cdots s_k \mid s_i \in S, \text{distinct}, k \leq n\}$, where $n = |S|$. In other words, \mathcal{S} is the set of all possible products of distinct generators. We prove that for any element $t \in W \setminus \mathcal{H}$, there exists $g \in \mathcal{S}$ such that $gt \in \mathcal{H}$.

Let $t \in W \setminus \mathcal{H}$ be given and consider any reduced expression \mathbf{t} for t . Multiply \mathbf{t} by all generators appearing an even (including zero) number of times in \mathbf{t} .

$$(s_{i_1} \cdots s_{i_n})\mathbf{t}$$

Then the resulting word, and hence any reduced expression, has the property that all generators appear and each generator appears an odd number of times. Therefore, the element represented by this word is essential, and hence, it has rank one.

$$W = \mathcal{H} \bigcup \left(\bigcup_{\alpha \in \mathcal{S}} \alpha \mathcal{H} \right).$$

This proves that $r(W) \leq 1$. ■

In order to prove that $\text{rank}(W) = 1$, we need to show $r(T) \leq 1$ for any finite index subgroup T of W . But it seems that the above argument does not work for T . Because, in the proof of Proposition 8, $(s_{i_1} \cdots s_{i_n})$ does not necessarily represent an element in T . Once one takes powers on $(s_{i_1} \cdots s_{i_n})$ to get an element in T , the argument fails to apply. In particular, if the index $[W : T]$ is even, all generators in $(s_{i_1} \cdots s_{i_n})^{[W:T]}$ appear an even number of times. Therefore, we take a different approach to prove $r(T) \leq 1$.

Definition 9 Let \mathbf{w} be a reduced word in S . For any generator s appearing in \mathbf{w} , let

$$\mathbf{w} = \mathbf{w}_0 s \mathbf{w}_1 s \cdots s \mathbf{w}_k s \mathbf{w}_{k+1},$$

where \mathbf{w}_i does not contain s for all $0 \leq i \leq k+1$. Note that \mathbf{w}_0 and \mathbf{w}_{k+1} are allowed to be empty, but each $\mathbf{w}_i \neq \emptyset$ for $1 \leq i \leq k$.

1. \mathbf{w} is said to be *s-minimal* if each subword \mathbf{w}_i , for $1 \leq i \leq k$, contains a *s-blocker*, i.e., a generator $s' \in S$ such that $ss' \neq s's$. We consider \mathbf{w} to be vacuously *s-minimal* if s appears only once in \mathbf{w} .
2. \mathbf{w} is said to be *s-good* if \mathbf{w} is *s-minimal* and $\mathbf{w}_{k+1}\mathbf{w}_0$ contains a *s-blocker* for $k \geq 1$. In the case that $k = 0$, \mathbf{w} is considered to be *s-good*.

Remark 10 Any reduced word \mathbf{w} is *s-minimal* for all generators s appearing in \mathbf{w} . For a generator s appearing in \mathbf{w} , let

$$\mathbf{w} = \mathbf{w}_0 s \mathbf{w}_1 s \cdots s \mathbf{w}_k s \mathbf{w}_{k+1}.$$

If \mathbf{w}_i does not contain a s -blocker for some $1 \leq i \leq k$, then $s\mathbf{w}_i s = \mathbf{w}_i$, which is a contradiction.

Lemma 11 *Let \mathbf{w} be a reduced word which is s -good for all $s \in S$. Then the element represented by \mathbf{w} is essential.*

Proof. Suppose that s appears once in \mathbf{w} . In other words, $\mathbf{w} = \mathbf{w}_0 s \mathbf{w}_1$. Suppose the element represented by \mathbf{w} is in $u^{-1}W_J u$ for some $u \in W$ and $J \subset S$. Then the element represented by $\mathbf{u}\mathbf{w}\mathbf{u}^{-1}$ lies in W_J , where \mathbf{u} is any reduced expression for u . In some (any) reduced expression of $\mathbf{u}\mathbf{w}\mathbf{u}^{-1}$, s appears an odd number of times. In particular, s appears. Therefore, $s \in J$.

Suppose s appears at least twice in \mathbf{w} . In other words,

$$\mathbf{w} = \mathbf{w}_0 s \mathbf{w}_1 s \cdots s \mathbf{w}_k s \mathbf{w}_{k+1},$$

for $k \geq 1$. Suppose that the element represented by \mathbf{w} is in $u^{-1}W_J u$ for some $u \in W$ and $J \subset S$. Then the element represented by

$$\mathbf{u}\mathbf{w}\mathbf{u}^{-1} = \mathbf{u}\mathbf{w}_0 s \mathbf{w}_1 s \cdots s \mathbf{w}_k s \mathbf{w}_{k+1} \mathbf{u}^{-1}$$

lies in W_J , where \mathbf{u} is any reduced expression of u .

Assume that $s \notin J$. Then \mathbf{u} must contain s . We prove that one of two s 's in \mathbf{u} and \mathbf{u}^{-1} cannot be cancelled off. By way of contradiction, let us assume that both can be cancelled.

Since \mathbf{w} is s -good, there exists at least one s -blocker in \mathbf{w}_0 or \mathbf{w}_{k+1} . Without loss of generality, we assume that s -blocker lies in \mathbf{w}_0 . (A symmetric argument applies if it lies in \mathbf{w}_{k+1} .) Take the first s -blocker in \mathbf{w}_0 and call it s_1 . It follows that \mathbf{u} must contain s_1 and the last s_1 occurs after the last s in \mathbf{u} .

$$\mathbf{u}\mathbf{w}\mathbf{u}^{-1} = (\cdots s \cdots s_1 \cdots)(\cdots s_1 \cdots) s \mathbf{w}_1 s \cdots s \mathbf{w}_k s \mathbf{w}_{k+1} (\cdots s_1 \cdots s \cdots)$$

In order for the last s to be cancelled off, s_1 must occur in \mathbf{w}_{k+1} .

$$\mathbf{u}\mathbf{w}\mathbf{u}^{-1} = (\cdots s \cdots s_1 \cdots)(\cdots s_1 \cdots) s \mathbf{w}_1 s \cdots s \mathbf{w}_k s (\cdots s_1 \cdots)(\cdots s_1 \cdots s \cdots),$$

where the $s_1 \in \mathbf{w}_{k+1}$ written above is the last occurrence of s_1 in \mathbf{w}_{k+1} . So before being able to cancel the first and the last s , we need to cancel out the intermediate blocker s_1 .

Now \mathbf{w} is s_1 -good. Therefore, there exists an s_1 -blocker on the left of the first $s_1 \in \mathbf{w}_0$ or on the right of the last $s_1 \in \mathbf{w}_{k+1}$. Take the first s_1 -blocker in \mathbf{w}_0 or the last s_1 -blocker in \mathbf{w}_{k+1} and call it s_2 . Note that $s_2 \neq s$ and before canceling the s_1 , we must first be able to cancel out the s_1 -blocker s_2 . As in the last paragraph, this forces $\mathbf{u}\mathbf{w}\mathbf{u}^{-1}$ to be of the form

$$(\cdots s_1 \cdots s_2 \cdots)(\cdots s_2 \cdots s_1 \cdots) s \mathbf{w}_1 s \cdots s \mathbf{w}_k s (\cdots s_1 \cdots s_2 \cdots)(\cdots s_2 \cdots s_1 \cdots)$$

Note that \mathbf{w} is s_2 -good, and hence, there exists an s_2 -blocker ($\neq s, s_1$) on the left of the first s_2 or on the right of the last s_2 in \mathbf{w} . But since $|S| < \infty$, this process must stop in finitely many stages, which proves that one of s 's in \mathbf{u} and \mathbf{u}^{-1} cannot be cancelled off. Therefore, $s \in J$. The element represented by \mathbf{w} is essential. \blacksquare

Let T be a proper finite index subgroup of W . Assume that T is normal and let $n = [W : T] \geq 2$. In order to prove that $r(T) \leq 1$, we need to consider two types of generators for a given reduced word \mathbf{w} representing an element in T : (1) a generator s does not appear in \mathbf{w} and (2) a generator s appears, but \mathbf{w} is not s -good. We begin with generators of type (1).

Let \mathbf{w} be a reduced word in S representing an element in T and assume that \mathbf{w} misses a generator s . Choose $s' \in S$ such that $ss' \neq s's$. Note that such s' always exists because W is infinite irreducible. Next choose $s'' \in S$ such that either $ss'' \neq s''s$ or $s's'' \neq s''s'$. Note that such s'' always exists, otherwise $W = W_{\{s,s'\}} \times W_{S \setminus \{s,s'\}}$, contradicting irreducibility (if $|S| > 2$) or non-affine (if $|S| = 2$).

Lemma 12 1. If $ss'' \neq s''s$, then $\mathbf{r} = (s''ss')^n \mathbf{w}$ has the following property:

- (a) The element represented by \mathbf{r} is in T . This is obvious.
- (b) \mathbf{r} is s -good.
- (c) \mathbf{r} is s' -good.
- (d) For $t \neq s, s', s''$, if \mathbf{w} is t -good, then \mathbf{r} is also t -good.

2. If $s's'' \neq s''s'$, then $\mathbf{r}' = (s's''ss')^n \mathbf{w}$ has the following property:

- (a) The element represented by \mathbf{r}' is in T .
- (b) \mathbf{r}' is s -good.
- (c) \mathbf{r}' is s'' -good.
- (d) For $t \neq s, s', s''$, if \mathbf{w} is t -good, then \mathbf{r}' is also t -good.

Proof. We prove the first statement only. The second statement can be proved by exactly the same argument as the first.

By hypothesis, $ss' \neq s's$, $ss'' \neq s''s$, and s does not appear in \mathbf{w} . Then

$$\mathbf{r} = (s''ss')^n \mathbf{w} = (s''ss')(s''ss') \cdots (s''ss') \mathbf{w} = (s'')s(s's'')s(s' \cdots s'')s(s'\mathbf{w})$$

is s -good as both s' and s'' are s -blockers. Let $\mathbf{w} = \mathbf{w}'_0 s' \mathbf{w}'_1 s' \cdots \mathbf{w}'_l s' \mathbf{w}'_{l+1}$. Then

$$\mathbf{r} = (s''ss')^n \mathbf{w} = (s''s)s'(s''s)s' \cdots (s''s)s' \mathbf{w}'_0 s' \mathbf{w}'_1 s' \cdots \mathbf{w}'_l s' \mathbf{w}'_{l+1}$$

Note that if the subword \mathbf{w}'_0 does not contain s' -blocker, then $s' \mathbf{w}'_0 s'$ is reduced to \mathbf{w}'_0 . But \mathbf{w}'_1 contains a s' -blocker and s is an s' -blocker. Therefore, there exists at least one s' -blocker in the reduced expression of $(s''ss') \mathbf{w}'_0 s' \mathbf{w}'_1$, even if \mathbf{w}'_0 does not contain s' -blocker. (Recall that the letter s does not appear in \mathbf{w}'_0 and \mathbf{w}'_1 , so it cannot be cancelled off.)

Suppose that \mathbf{w} is t -good for $t \neq s, s', s''$. If neither s, s' or s'' is t -blocker, then \mathbf{r} is obviously t -good. Consider the case that the t -blocker is either s, s' or s'' .

$$\mathbf{r} = (s''ss')^n \mathbf{w} = (s''ss')(s''ss') \cdots (s''ss') \mathbf{w}'_0 t \mathbf{w}'_1 t \cdots \mathbf{w}'_m t \mathbf{w}'_{m+1}$$

Note that s is not in \mathbf{w} and $n \geq 2$. Therefore, there exists at least one t -blocker in the reduced expression of $(s''ss')(s''ss') \cdots (s''ss') \mathbf{w}'_0$. It follows that \mathbf{r} is t -good. \blacksquare

- Remark 13** 1. In Lemma 12, \mathbf{r} is not necessarily s'' -good. Similarly, \mathbf{r}' is not necessarily s' -good. Therefore, the number of good generators of \mathbf{r} or \mathbf{r}' might be equal to the number of good generators of \mathbf{w} . But note that s appears in \mathbf{r} and \mathbf{r}' , and all letters appearing in \mathbf{w} still appear in \mathbf{r} and \mathbf{r}' .
2. Let \mathbf{w} be a reduced word in S representing an element in T . By multiplying words as in Lemma 12, we can obtain $\mathbf{w}' := \mathbf{y}_k \mathbf{y}_{k-1} \cdots \mathbf{y}_1 \mathbf{w}$ such that the element represented by \mathbf{w}' is in T and all generators appear in \mathbf{w}' .
3. There exists a finite set \mathcal{R} of words such that for any reduced word \mathbf{w} representing an element T , there exists some $\mathbf{r} \in \mathcal{R}$ for which $\mathbf{r}\mathbf{w}$ represents an element in T and all generators appear in $\mathbf{r}\mathbf{w}$.

Next, we consider generators of type (2).

Definition 14 Let \mathbf{w} be a reduced word in S such that all generators appear. Define $B(\mathbf{w})$ be the set of generators for which \mathbf{w} is not good. For $B \subset S$, a word \mathbf{v} is called a B -cancellator if, for any \mathbf{w} such that $B = B(\mathbf{w})$, $\mathbf{v}\mathbf{w}$ is s -good for all $s \in S$.

The following lemma tells us that B -cancellators exist and can be chosen to represent an element in T . Note that there are only finitely many subsets of S . Therefore, we can form finitely many cancellators.

Let \mathbf{w} be a reduced word in S such that all generators appear and \mathbf{w} represents an element in T . Suppose that \mathbf{w} is not s -good. Choose $s' \in S$ such that $ss' \neq s's$. Note that such s' always exists because W is infinite irreducible. Also we choose $s'' \in S$ such that either $ss'' \neq s''s$ or $s's'' \neq s''s'$. Note that such s'' always exists, otherwise $W = W_{\{s, s'\}} \times W_{S \setminus \{s, s'\}}$.

Lemma 15 1. If $ss'' \neq s''s$, then $\mathbf{r} = (s''ss')^n \mathbf{w}$ has the following property:

- (a) The element represented by \mathbf{r} is in T . This is obvious.
- (b) \mathbf{r} is s -good.
- (c) \mathbf{r} is s' -good.
- (d) \mathbf{r} is s'' -good.
- (e) For $t \neq s, s', s''$, if \mathbf{w} is t -good, then \mathbf{r} is also t -good.

2. If $s's'' \neq s''s'$, then $\mathbf{r}' = (s's''ss's'')^n \mathbf{w}$ has the following property:

- (a) The element represented by \mathbf{r}' is in T .
- (b) \mathbf{r}' is s -good.
- (c) \mathbf{r}' is s' -good.
- (d) \mathbf{r}' is s'' -good.
- (e) For $t \neq s, s', s''$, if \mathbf{w} is t -good, then \mathbf{r}' is also t -good.

Proof. Again, we prove the first statement only. The second statement can be proved by exactly the same argument as the first.

Let

$$\mathbf{r} = (s''ss')^n \mathbf{w} = (s'')s(s's'') \cdots (s's'')s(s'\mathbf{w}_0)s\mathbf{w}_1s \cdots s\mathbf{w}_ks\mathbf{w}_{k+1}.$$

In the reduced expression of $ss'\mathbf{w}_0s$, since s and s' don't commute, s' is s -blocker. (Note that \mathbf{w}_0 does not contain s' , since, by assumption, $s \in B(\mathbf{w})$). Also the first s'' is an s -blocker. Hence \mathbf{r} is s -good.

$$\mathbf{r} = (s''ss')^n \mathbf{w} = (s'')s'(s''s)s' \cdots (s''s)s'\mathbf{w}'_0s'\mathbf{w}'_1s' \cdots \mathbf{w}'_ms'\mathbf{w}'_{j+1}$$

Secondly, note that s and s' don't commute. Since \mathbf{w} is not s -good, the first s' in \mathbf{w} should occur after the first s in \mathbf{w} , i.e., $s \in \mathbf{w}'_0$. It follows that s is an s' -blocker in \mathbf{w}'_0 . Hence \mathbf{r} is s' -good.

$$\mathbf{r} = (s''ss')^n \mathbf{w} = s''(ss')s''(ss') \cdots s''(ss'\mathbf{w}''_0)s''\mathbf{w}''_1s'' \cdots \mathbf{w}''_is''\mathbf{w}''_{i+1}$$

Note that s and s'' don't commute. Since \mathbf{w} is not s -good, the last s'' in \mathbf{w} should occur before the last s in \mathbf{w} , i.e., $s \in \mathbf{w}'_{i+1}$. It follows that \mathbf{w}''_{i+1} contains an s'' -blocker s . \mathbf{r} is s'' -good.

Suppose that \mathbf{w} is t -good for $t \neq s, s', s''$. If neither s, s' or s'' is t -blocker, then \mathbf{r} is t -good. Let

$$\mathbf{r} = (s''ss')^n \mathbf{w} = (s''ss')(s''ss') \cdots (s''ss')\mathbf{w}'_0t\mathbf{w}'_1t \cdots \mathbf{w}'_ht\mathbf{w}'_{h+1}.$$

If s is a t -blocker, the first t occur after the first s in \mathbf{w} , i.e., $s \in \mathbf{w}'_0$. Furthermore, this s cannot be cancelled off, because s and s' don't commute. It follows that there exists at least one t -blocker in the reduced expression of $(s''ss')\mathbf{w}'_0$. \mathbf{r} is t -good. Next, suppose that s' or s'' is t -blocker. The s in the last $(s''ss')$ cannot be cancelled off. Therefore, there exists at least one t -blocker in the reduced expression of $(s''ss')(s''ss') \cdots (s''ss')\mathbf{w}'_0$. \mathbf{r} is t -good. ■

Corollary 16 *For a given $B \subset S$, there exists a B -cancellator \mathbf{v}_B and it represents an element in T .*

Proof. Let \mathbf{w} be a reduced word in S , with all generators appearing, such that $B(\mathbf{w}) = B$. Choose a generator $s_1 \in B(\mathbf{w})$. Apply the lemma to obtain a word $\mathbf{v}_1 \in T$ such that $\mathbf{v}_1\mathbf{w}$ is s_1 -good. Consider $\mathbf{v}_1\mathbf{w}$. From Lemma 15, $B(\mathbf{v}_1\mathbf{w}) \subset B(\mathbf{w})$ and $|B(\mathbf{v}_1\mathbf{w})| < |B(\mathbf{w})|$. Choose a generator $s_2 \in B(\mathbf{v}_1\mathbf{w})$ and apply the lemma to obtain a word $\mathbf{v}_2 \in T$ such that $\mathbf{v}_2\mathbf{v}_1\mathbf{w}$ is s_2 -good. Continuing this process, at most $|B(\mathbf{w})|$ number of times, we obtain a word $\mathbf{v}_k\mathbf{v}_{k-1} \cdots \mathbf{v}_1\mathbf{w}, k \leq |B(\mathbf{w})|$ which is s -good for all $s \in S$. Such a $\mathbf{v}_k\mathbf{v}_{k-1} \cdots \mathbf{v}_1$ is the desired B -cancellator. ■

Corollary 17 $r(T) \leq 1$.

Proof. Let $\mathcal{H}_T = \mathcal{H} \cap T$. For $g \in \mathcal{H}_T$, the centralizer $C_W(g)$ in W is virtually infinite cyclic, and hence, $C_T(g)$ is also virtually infinite cyclic. It follows that $\mathcal{H}_T \subset \mathcal{A}_1(T)$.

Let $g' \in T \setminus \mathcal{H}_T$ be given. By Remark 13 and Corollary 16, we can find a $\mathbf{r} \in \mathcal{R}$ and a $B(\mathbf{r}\mathbf{w}')$ -cancellator \mathbf{v} , where \mathbf{w}' is any reduced expression for g' , such that $\mathbf{v}\mathbf{r}\mathbf{w}'$ is s -good for all $s \in S$, and hence, $\mathbf{v}\mathbf{r}\mathbf{w}'$ represents an element in \mathcal{H}_T . Therefore,

$$T = \bigcup_{B \subseteq S, \mathbf{r} \in \mathcal{R}} \mathbf{r}^{-1} \mathbf{v}_B^{-1} \mathcal{H}_T,$$

where each \mathbf{v}_B is a B -cancellator. \blacksquare

Proposition 18 *Let W be an infinite, irreducible, and non-affine right-angled Coxeter group. Then $\text{rank}(W) = 1$.*

Proof. By Proposition 8, it suffices to show that $r(W') \leq 1$ for any finite index subgroup W' of W . By taking the normal core, we obtain a finite index normal subgroup W'' of W and, by Corollary 17, $r(W'') \leq 1$. Finally, Proposition 4 implies that $r(W') \leq 1$. \blacksquare

We close the subsection by introducing two corollaries of Proposition 18. Two groups G_1 and G_2 are said to be *commensurable* if there exist H_i , for $i = 1, 2$, such that $[G_i : H_i] < \infty$ or $[H_i : G_i] < \infty$ and H_1 is isomorphic to H_2 . Davis proved that any infinite irreducible non-affine Coxeter group W cannot be a uniform lattice Γ in a higher rank non-compact connected semi-simple Lie group (See [8, Corollary 10.9.8]) and conjectured W cannot be commensurable to Γ . (See [7]) The conjecture is known to be true, see for example, [6], [14], [27]. Proposition 18 provides a new proof of the conjecture for right-angled Coxeter groups.

Corollary 19 *Let W be an infinite irreducible non-affine right-angled Coxeter group. Then W is not commensurable to any uniform lattice in a higher rank non-compact connected semi-simple Lie group G .*

Proof. Let Λ be a uniform lattice in G . By [2, Theorem 3.11], $\text{rank}(\Lambda) = \text{rank}(G) \geq 2$. Applying Proposition 4, we obtain that any group Γ commensurable to Λ satisfies $\text{rank}(\Gamma) = \text{rank}(\Lambda) \geq 2$. On the other hand, for any finite index subgroup W' of W , $\text{rank}(W) = \text{rank}(W') = 1$. Therefore, W and Λ cannot be commensurable. \blacksquare

Two metric spaces X and Y are said to be *quasi-isometric* if there exists a map $f : X \rightarrow Y$ such that, for some $\lambda \geq 1, \epsilon \geq 0$ and $C \geq 0$, such that

1. For all $x_1, x_2 \in X$,

$$\frac{1}{\lambda} d_X(x_1, x_2) - \epsilon \leq d_Y(f(x_1), f(x_2)) \leq \lambda d_X(x_1, x_2) + \epsilon.$$

2. Y lies in the C -neighborhood of the image of f .

The other corollary follows from Quasi-Isometry rigidity theorem due to Kleiner-Leeb ([19]) and Eskin-Farb ([12]).

Corollary 20 *Let W be an infinite irreducible non-affine right-angled Coxeter group. Then W is not quasi-isometric to any uniform lattice in a higher rank non-compact connected semi-simple Lie group G .*

Proof. By QI-rigidity theorem, if W is quasi-isometric to a uniform lattice in a higher rank non-compact connected semi-simple Lie group, W should be commensurable to a lattice. By Corollary 19, W cannot be commensurable to the lattice. ■

3.2 Algebraic Ranks of Right-Angled Artin Groups

An Artin group A is a group with the following presentation :


$$A = \langle s_1, \dots, s_n \mid \underbrace{s_i s_j \cdots}_{m_{ij}} = \underbrace{s_j s_i \cdots}_{m_{ij}}, \text{ for all } i \neq j \rangle,$$

where $m_{ij} = m_{ji}$ is an integer ≥ 2 or $m_{ij} = \infty$ in which case we omit the relation between s_i and s_j . As one can see, by adding relations $s_i = s_i^{-1}$ to the presentation, we obtain a Coxeter group. Right-angled Artin groups are those Artin groups for which all $m_{ij} = 2$ or ∞ for $i \neq j$.

One of the easy ways of defining a right-angled Coxeter group or a right-angled Artin group is via *the defining graph* Γ . This is the graph whose vertices are labeled by $S = \{s_1, \dots, s_n\}$ and two vertices s_i and s_j are connected if $m_{ij} = 2$. We denote by A_Γ (W_Γ , respectively) the right-angled Artin group (the right-angled Coxeter group, respectively) associated to a finite simplicial graph Γ . For example, if Γ consists of n vertices and no edges, then A_Γ is the free group on n generators. At the other extreme, if Γ is a complete graph with n vertices, A_Γ is the free abelian group of rank n .

Analogous to the Coxeter group situation, there is a CAT(0) space associated to a right-angled Artin group A_Γ , which can be constructed by the following process : begin with a wedge of circles attached to a point x_0 and labeled by the generators s_1, \dots, s_n . For each edge connecting s_i and s_j in Γ , attach a 2-torus with boundary labeled by the relator $s_i s_j s_i^{-1} s_j^{-1}$. For each triangle connecting s_i, s_j, s_k in Γ , attach a 3-torus with faces corresponding to the tori for the three edges of triangle. Continuing this process, attach a k -torus for each set of k -mutually commuting generators. The resulting cube complex is called *a Salvetti complex* for A_Γ and denoted by \mathcal{S}_Γ . It is easy to verify that the fundamental group of \mathcal{S}_Γ is A_Γ and the link of the unique vertex x_0 is a flag. It follows from Gromov's criterion that the universal cover X_Γ of the complex \mathcal{S}_Γ is a CAT(0) cube complex, and A_Γ acts on X_Γ freely and cocompactly.

Given two graphs Γ_1, Γ_2 , their *join* is the graph obtained by connecting every vertex of Γ_1 to every vertex of Γ_2 . If Γ is the join of Γ_1 and Γ_2 , then $A_\Gamma = A_{\Gamma_1} \times A_{\Gamma_2}$ and $X_\Gamma = X_{\Gamma_1} \times X_{\Gamma_2}$. We prove



Proposition 21 *If Γ is not a join, then $\text{rank}(A_\Gamma) = 1$.*

Remark 22 If Γ is the join of Γ_1 and Γ_2 , then

$$\text{rank}(A_\Gamma) = \text{rank}(A_{\Gamma_1}) + \text{rank}(A_{\Gamma_2}) \geq 2.$$

The proof of Proposition 21 is a direct consequence of a theorem due to Davis and Januszkiewicz ([9]). For a given graph Γ , we define two graphs Γ' and Γ'' as follows : The vertex set of Γ'' is $I \times \{0, 1\}$, where I is the vertex set of Γ . Two vertices $(i, 1)$ and $(j, 1)$ in $I \times 1$ are connected by an edge in Γ'' if and only if the corresponding vertices i and j span an edge in Γ . Any two distinct vertices in $I \times 0$ are connected by an edge. Finally, vertices $(i, 0)$ and $(j, 1)$ are connected by an edge if and only if $i \neq j$. The vertex set of Γ' is $I \times \{-1, 1\}$. The subsets $I \times (-1)$ and $I \times 1$ both span copies of Γ . A vertex $(i, -1)$ in $I \times (-1)$ is connected to $(j, 1)$ in $I \times 1$ if and only if $i \neq j$ and the vertices i and j span an edge of Γ . (See above for an example.) Then

Theorem 23 [9] *A_Γ and $W_{\Gamma'}$ are subgroups of $W_{\Gamma''}$ of index 2^I .*

Lemma 24 *Γ is a join if and only if Γ' is a join.*

Proof. For any subset $I' \subset I$, let $\Gamma_{I'}$ be a full subgraph of Γ whose vertex set is I' . It is obvious that if Γ is a join, then Γ' is a join. Namely, if Γ is a join of Γ_{I_1} and Γ_{I_2} , then Γ' is a join of $\Gamma'_{I_1 \times \{-1, 1\}}$ and $\Gamma'_{I_2 \times \{-1, 1\}}$. Conversely, suppose Γ' is a join of $\Gamma'_{I'_1}$ and $\Gamma'_{I'_2}$. Note that a vertex $(i, 1) \in \Gamma'_{I'_1}$ if and only if $(i, -1) \in \Gamma'_{I'_1}$. Therefore, Γ' is a join of $\Gamma'_{I_1 \times \{-1, 1\}}$ and $\Gamma'_{I_2 \times \{-1, 1\}}$ for some $I_1, I_2 \subset I$ and Γ is a join of Γ_{I_1} and Γ_{I_2} . ■

Proof of Proposition 21. Suppose that Γ is not a join. By Lemma 24, the corresponding graph Γ' is not a join. It follows that $W_{\Gamma'}$ is irreducible. If $W_{\Gamma'}$ has a finite index free abelian subgroup K , then $K \cap A_\Gamma$ is also a finite index free abelian subgroup of A_Γ . Since Γ is not a join, this is impossible. By Proposition 18, $\text{rank}(W_{\Gamma'}) = 1$, and hence, $\text{rank}(A_\Gamma) = 1$. ■

Corollary 25 *Let A be a right-angled Artin group which is not a join. Then A is not commensurable (or quasi-isometric) to any uniform lattice in a non-compact connected semi-simple Lie group of higher rank.*

4 Relatively Hyperbolic Groups

Relatively hyperbolic groups are a generalization of hyperbolic groups. They were introduced by Gromov ([15]) and many equivalent definitions have been developed by different authors in different contexts. See, for example, Farb([13]), Bowditch([3]), Yaman([30]), Druţu-Osin-Sapir([11]), Osin([24]), Druţu([10]), and Mineyev-Yaman([22]). In this paper, we shall use the Bowditch's definition via geometrically finite convergence groups.

Suppose that M is a compact metrizable topological space and a group G acts by homeomorphisms on M . G is said to be a *convergence group* if the induced action on the space of distinct triples is properly discontinuous. Let G be a convergence group on M . A point $\xi \in M$ is a *conical limit point* if there exists a sequence, $\{g_n\}$, in G and two points $\zeta \neq \eta \in M$, such that $g_n(\xi) \rightarrow \zeta$ and $g_n(\xi') \rightarrow \eta$ for all $\xi' \neq \xi$. An element $g \in G$ is a *hyperbolic element* if it is of infinite order and fixes exactly two points in M . A subgroup $H < G$ is *parabolic* if H is infinite, fixes a point $\xi \in M$, and contains no hyperbolic elements. In this case, the fixed point of H is unique and we call the point a *parabolic point*. The nontrivial elements in a parabolic subgroup are called *parabolic elements*. If ζ is a parabolic point, the stabilizer of ζ , $\text{Stab}(\zeta)$, is necessarily a parabolic subgroup. A parabolic point ζ is a *bounded parabolic point* if the stabilizer $\text{Stab}(\zeta)$ acts properly and cocompactly on $M \setminus \{\zeta\}$. Finally, a convergence group G on M is a *geometrically finite group* if each point of M is either a conical limit point or a bounded parabolic point.

Definition 26 A group G is *hyperbolic relative to a family of infinite finitely generated subgroups \mathcal{G}* if it acts properly discontinuously by isometries on a proper geodesic hyperbolic space X such that the induced action on ∂X is of convergence, geometrically finite, and such that the maximal parabolic subgroups are exactly the elements of \mathcal{G} . Elements of \mathcal{G} are called *peripheral subgroups*.

It is known that all definitions mentioned above are equivalent, provided that the group G and all peripheral subgroups are infinite and finitely generated. But some authors do not assume that peripheral subgroups are infinite and finitely generated. In fact, it has been shown in [30] that the finite generation of peripheral subgroups can be dispensed with. Also some definitions allow the elements of \mathcal{G} to be finite. But, in [24], Osin proved that one can make \mathcal{G} smaller so that all peripheral subgroups are infinite (or possibly empty). The followings are well-known examples of relatively hyperbolic groups.

Example 27 • Hyperbolic groups: These are hyperbolic relative to $\mathcal{G} = \emptyset$.

- Geometrically finite isometry groups of Hadamard manifolds of negatively pinched sectional curvature: These are hyperbolic relative to the maximal parabolic subgroups.
- Free products of finitely many finitely generated groups: These are hyperbolic relative to the factors, since the action on the Bass-Serre tree satisfies the second definition of Bowditch. See [3, Definition 2].

- Groups G acting geometrically on a CAT(0) space X which has the isolated flats property: In this case, X is an asymptotically tree-graded space and G is hyperbolic relative to the collection of virtually abelian subgroups of rank at least two. (See [17])

In the next subsection, we will prove that if G is a relatively hyperbolic group with $|\mathcal{G}| \geq 2$, and there are two peripheral subgroups containing elements of infinite order, then $\text{rank}(G) \leq 1$. The following theorem of Bowditch on the existence of an invariant collection of disjoint horoballs provides the crucial tool in our proof.

Let X be a δ -hyperbolic geodesic metric space for some $\delta > 0$ and $\xi \in \partial X$. A function $h : X \rightarrow \mathbb{R}$ is a *horofunction about ξ* if there exist constants $c_1 = c_1(\delta), c_2 = c_2(\delta)$ such that if $x, a \in X$ and $d(a, x\xi) \leq c_1$, for some geodesic ray $x\xi$ from x to ξ , then $|h(a) - h(x) - d(x, a)| \leq c_2$. A closed set $B \subset X$ is a *horoball about ξ* if there is a horofunction h about ξ and a constant $c = c(\delta)$ such that $h(x) \geq -c$ for all $x \in B$, and $h(x) \leq c$ for all $x \in X \setminus B$. In this case ξ is called the center of the horoball and is uniquely determined by B .

Proposition 28 [3, Proposition 6.13] *Let G be a relatively hyperbolic group and X a space on which G acts as in Definition 26. Let Π be the set of all bounded parabolic points in ∂X . Then Π/G is finite. Moreover, for any $r > 0$, there is a collection of horoballs $\mathcal{B} = \{B_\xi | \xi \in \Pi\}$ indexed by Π with the following properties*

1. \mathcal{B} is r -separated, that is, $d(B_\xi, B_\eta) \geq r$ for all $\xi \neq \eta \in \Pi$.
2. \mathcal{B} is G -invariant, that is, $g(B_\xi) = B_{g(\xi)}$ for all $g \in G$ and $\xi \in \Pi$.
3. $Y(\mathcal{B})/G$ is compact, where $Y(\mathcal{B}) = X \setminus \bigcup_{\xi \in \Pi} \text{int}(B_\xi)$.

Note that the intersection of any two peripheral subgroups is finite and there are finitely many conjugacy classes of peripheral subgroups.

4.1 Algebraic Ranks of Relatively Hyperbolic Groups

In order to prove that a group has an algebraic rank ≤ 1 , we need to figure out the set $\mathcal{A}_1(G)$ and show that the group can be covered by finitely many translates of $\mathcal{A}_1(G)$. Also the procedure needs to be repeated for all finite index subgroups. We introduce two lemmas which enable us to find the elements by which the group can be covered by translates of $\mathcal{A}_1(G)$.

For a finite set of isometries F of a metric space X and $x \in X$, let $\lambda(x, F) = \max\{d(f(x), x) | f \in F\}$.

Lemma 29 [20] *Let X be a δ -hyperbolic geodesic metric space and G a group of isometries of X with a finite generating set S . If $\lambda(x, S) > 100\delta$ for all $x \in X$, then G contains a hyperbolic element g such that $d_S(\text{id}, g) = 1$ or 2.*

Hereafter, suppose that G is a relatively hyperbolic group with $|\mathcal{G}| \geq 2$ and X a proper δ -hyperbolic geodesic space on which G acts as in Definition 26. Also we assume that there are at least two peripheral subgroups in \mathcal{G} containing elements of

infinite order. Lemma 28 implies that there is a 200δ -separated invariant collection of horoballs \mathcal{B} centered at the parabolic points such that $Y(\mathcal{B})/G$ is compact.

Lemma 30 [29, Lemma 3.1] *There exists a positive integer k_1 with the following property : for any infinite order element $\gamma \in G$ and any $x \in Y(\mathcal{B})$, there is some $k, 1 \leq k \leq k_1$, such that $d(\gamma^k(x), x) \geq 200\delta$.*

Let \mathcal{H} be the set of hyperbolic elements.

Proposition 31 $\mathcal{H} \subset \mathcal{A}_1(G)$.

Proof. Let $g \in \mathcal{H}$ be given and A the two fixed points of $\langle g \rangle$ in ∂X . If $h \in G$ commutes with g , then h fixes A (See [28, Corollary 2O]). Combining this with the fact that $\langle g \rangle$ is of finite index in the stabilizer $H = \{k \in G \mid kA = A\}$ (see [28, Theorem 2I]), the centralizer of g in G , $C_G(g)$ has a free abelian group of rank at most one as a finite index subgroup. Therefore, $\mathcal{H} \subset \mathcal{A}_1(G)$. ■

Choose two elements of infinite order from two different peripheral subgroups and denote them by h_1 and h_2 . We also denote the horoball stabilized by h_i by $B_i, i = 1, 2$. Since h_i is chosen to be of infinite order, B_i is the only horoball stabilized by $h_i, i = 1, 2$. Then

Proposition 32 *Let $g \in G \setminus \mathcal{H}$ be an infinite parabolic element. Then $h_i^k g$ is hyperbolic for some $i = 1, 2$ and for some $k, 1 \leq k \leq k_1$, where k_1 is the constant appearing in Lemma 30.*

Proof. Suppose that g is infinite parabolic. Without loss of generality, h_1 and g are contained in different peripheral subgroups. Following Lemma 30, consider $K = \langle h_1^k, g \rangle$. We prove that $\lambda(x, \{h_1^k, g\}) > 100\delta$ for any $x \in X$. Suppose that there exists some $x \in B$ for some $B \in \mathcal{B}$ such that $\lambda(x, \{h_1^k, g\}) \leq 100\delta$. Then $h_1^k(B) = B$ and $g(B) = B$ (Note that \mathcal{B} is 200δ -separated). It follows that the center of B is fixed by K . Since h_1^k and g fix two different centers, this is a contradiction. By Lemma 30, for any $x \in Y(\mathcal{B})$, there is a $1 \leq k \leq k_1$ such that $d(h_1^k(x), x) \geq 200\delta$. Therefore, $\lambda(x, \{h_1^k, g\}) \geq 200\delta$. Lemma 29 implies that $h_1^k g$ or $g h_1^k$ is hyperbolic. (note that both h_1^k and g are parabolic.) Furthermore, if $g h_1^k$ is hyperbolic, then $h_1^k g$ is also hyperbolic. In fact, suppose that $g h_1^k$ is hyperbolic and fixes exactly two distinct points α and β , then $h_1^k g$ fixes $h_1^k(\alpha)$ and $h_1^k(\beta)$. Conversely, suppose $h_1^k g$ fixes a point $\gamma \neq h_1^k(\alpha), h_1^k(\beta)$. Choose γ' such that $h_1^k(\gamma') = \gamma$. Note that $\gamma' \neq \alpha, \beta$. But $h_1^k g h_1^k(\gamma') = h_1^k(\gamma') \Rightarrow g h_1^k(\gamma') = \gamma'$. Therefore, $\gamma' = \alpha$ or β , which is a contradiction. ■

Proposition 33 $\text{rank}(G) \leq 1$

Proof. We decompose the set of torsion elements into $\mathcal{E} \amalg \mathcal{E}'$ as follows. A torsion element $g \in \mathcal{E}$ if and only if g stabilizes both B_1 and B_2 . Otherwise $g \in \mathcal{E}'$. Suppose that $g \in \mathcal{E}'$. Without loss of generality, assume that g does not stabilize B_1 . Then

we have $\lambda(x, \{h_1^k, g\}) \geq 100\delta$. In particular, $d(g(x), x) \geq 100\delta$ for $x \in B_1$. The same argument as in Proposition 32 implies that $h_1^k g$ is hyperbolic. Since the intersection of two distinct peripheral subgroups is at most finite, $|\mathcal{E}| < \infty$, say $\mathcal{E} = \{l_1, \dots, l_n\}$. Choose any hyperbolic element $h \in G$ and let $t_i = l_i h^{-1}$ for $i = 1, \dots, n$.

By combining with Proposition 32,

$$G = \mathcal{H} \bigcup \left(\bigcup_{i=1}^{k_1} h_1^{-i} \mathcal{H} \right) \bigcup \left(\bigcup_{i=1}^{k_1} h_2^{-i} \mathcal{H} \right) \bigcup \left(\bigcup_{i=1}^n t_i \mathcal{H} \right).$$

Therefore, $r(G) \leq 1$.

Next we need to prove that $r(T) \leq 1$ for any finite index subgroup T in G . By taking the normal core of T , it suffices to show that $r(T) \leq 1$ for any finite index normal subgroup T in G . Recall $r(G') \geq r(G)$ if G' is a finite index subgroup of G . Let T be a finite index normal subgroup in G and $m = [G : T]$. Also let $\mathcal{H}_T = \mathcal{H} \cap T$. Recall that \mathcal{H} is the set of hyperbolic elements in G . Then $\mathcal{H}_T \subset \mathcal{A}_1(T)$. It can be easily verified that all arguments in proving $r(G) \leq 1$ applies without any change to prove $r(T) \leq 1$, namely,

- $h_i^m \in T$ is an infinite parabolic element and stabilizes a horoball B_i for $i = 1, 2$.
- For any $g \in T \setminus \mathcal{H}_T$ of infinite order, $(h_i^m)^k g \in \mathcal{H}_T$ for some $i = 1, 2$.
- One can decompose the set of torsion elements in T as follows : $g \in \mathcal{E}_T$ if and only if g stabilizes both B_1 and B_2 . Otherwise $g \in \mathcal{E}'_T$. For any element g in \mathcal{E}'_T , $(h_i^m)^k g \in \mathcal{H}_T$ for some $i = 1, 2$. Since \mathcal{E}_T is finite, one can choose any hyperbolic element in T such that \mathcal{E}'_T can be covered by finitely many translates of \mathcal{H}_T .

■

4.2 CAT(0) Spaces with Isolated Flats

CAT(0) spaces with isolated flats were first introduced by Kapovich-Leeb and Wise, independently. In [18], Kapovich and Leeb study a class of CAT(0) spaces in which the maximal flats are disjoint and separated by regions of strictly negative curvature. Since then, they have been studied by a number of authors, in particular, because of their strong connections to relatively hyperbolic groups.

A *k-flat* is an isometrically embedded copy of Euclidean space \mathbb{E}^k for $k \geq 2$. A CAT(0) space X with a geometric group action (i.e., properly discontinuously and cocompactly by isometries) has *isolated flats* if it contains an equivariant collection \mathcal{F} of flats such that \mathcal{F} is closed and isolated in $\text{Flat}(X)$, and each flat $F \subset X$ in the space is contained in a uniformly bounded tubular neighborhood of some $F' \in \mathcal{F}$.

Let X be a CAT(0) space with isolated flats and G be a group acting geometrically on X . One of main results in [17] is

Theorem 34 [17, Theorem 1.2.1] *The following are equivalent.*

1. X has isolated flats.
2. Each component of the Tits boundary $\partial_T X$ is either an isolated point or a standard Euclidean sphere.
3. X is a relatively hyperbolic space with respect to a family of flats in \mathcal{F} .
4. G is a relatively hyperbolic group with respect to a collection of virtually abelian subgroups of rank at least two.

Proposition 35 *Let G be a group acting geometrically on a $\text{CAT}(0)$ space with isolated flats and $|\mathcal{F}| \geq 2$. Then $\text{rank}(G) \leq 1$.*

Proof. By Theorem 34, G is hyperbolic relative to a collection of virtually abelian subgroups of rank at least two. Since we assume that $|\mathcal{F}| \geq 2$, there are at least two peripheral subgroups in G . Proposition 33 applies that $\text{rank}(G) \leq 1$. ■

Remark 36 In the case that \mathcal{F} consists of a single flat F , one can conclude that G acts geometrically on F . Therefore, $\text{rank}(G)$ is equal to $\dim(F) \geq 2$ by [2].

References

- [1] W. Ballmann. Nonpositively curved manifolds of higher rank. *Ann. of Math.*, 122(3):597–609, 1985.
- [2] W. Ballmann and P. Eberlein. Fundamental groups of manifolds of nonpositive curvature. *J. Differential Geom.*, 25(1):1–22, 1987.
- [3] B. Bowditch. Relatively hyperbolic groups. *Internat. J. Algebra Comput.*, 22(3):1250016, 66pp, 2012.
- [4] K. Burns and R. Spatzier. Manifolds of nonpositive curvature and their buildings. *Inst. Hautes Études Sci. Publ. Math.*, pages 25–59, 1987.
- [5] P. Caprace and K. Fujiwara. Rank-one isometries of buildings and quasi-morphisms of kac-moody groups. *Geom. Funct. Anal.*, 19(5):1296–1319, 2010.
- [6] D. Cooper and D. D. Long. Infinite coxeter groups are virtually inducible. *Proc. Edinburgh Math. Soc. (2)*, 41(2):303–313, 1998.
- [7] M. Davis. Groups generated by reflections and aspherical manifolds not covered by euclidean space. *Ann. of Math.*, 117(2):293–324, 1983.
- [8] M. Davis. *The geometry and topology of Coxeter groups*. Princeton University Press, Princeton, 2008.
- [9] M. Davis and T. Januszkiewicz. Right-angled artin groups are commensurable with right-angled coxeter groups. *J. Pure Appl. Algebra*, 153(3):229–235, 2000.

- [10] C. Druţu. Relatively hyperbolic groups : geometry and quasi-isometric invariance. *Comment. Math. Helv.*, 84(3):503–546, 2009.
- [11] C. Druţu and M. Sapir. Tree-graded spaces and asymptotic cones of groups. *Topology*, 44(5):959–1058, 2005.
- [12] A. Eskin and B. Farb. Quasi-flats and rigidity in higher rank symmetric spaces. *J. Amer. Math. Soc.*, 10(3):653–692, 1997.
- [13] B. Farb. Relatively hyperbolic groups. *Geom. Funct. Anal.*, 8(5):810–840, 1998.
- [14] C. Gonciulea. Infinite coxeter groups virtually subject onto \mathbb{Z} .(english summary). *Comment. Math. Helv.*, 72(2):257–265, 1997.
- [15] M. Gromov. Hyperbolic groups. In *Essays in group theory*, volume 8 of *Math. Sci. Res. Inst. Publ.*, pages 75–263. Springer, New York, 1987.
- [16] P. Hall. Finiteness conditions for soluble groups. *Proc. London Math. Soc.* (3), 4:419–436, 1954.
- [17] G. Hruska and B. Kleiner. Hadamard spaces with isolated flats. *Geom. Topol.*, 9:1501–1538, 2005.
- [18] M. Kapovich and B. Leeb. On asymptotic cones and quasi-isometry classes of fundamental groups of 3-manifolds. *Geom. Funct. Anal.*, 5:582–603, 1995.
- [19] B. Kleiner and B. Leeb. Rigidity of quasi-isometries for symmetric spaces and euclidean buildings. *Inst. Hautes Etudes Sci. Publ. Math.*, 86:115–197, 1997.
- [20] M. Koubi. Croissance uniforme dans les groupes hyperboliques. *Ann. Inst. Fourier (Grenoble)*, 48(5):1441–1453, 1998.
- [21] D. Krammer. *The conjugacy problem for Coxeter groups*. PhD thesis, Universiteit Utrecht, 1994.
- [22] I. Mineyev and A. Yaman. Relative hyperbolicity and bounded cohomology, 2006. <http://www.math.uiuc.edu/~mineyev/math/>.
- [23] G. Moussong. *Hyperbolic Coxeter Groups*. PhD thesis, Ohio State, 1988.
- [24] D. V. Osin. Relatively hyperbolic groups : intrinsic geometry, algebraic properties, and algorithmic problems. *Mem. Amer. Math. Soc.*, 179(843):vi+100, 2006.
- [25] A. Ould Houcine. Embeddings in finitely presented groups which preserve the center. *J. Algebra*, 307(1):1–23, 2007.
- [26] G. Prasad and M. Raghunathan. Cartan subgroups and lattices in semi-simple groups. *Ann. of Math.*, 96:296–317, 1972.
- [27] S. Singh. Coxeter groups are not higher rank arithmetic groups., 2012. arXiv:1208.6569.

- [28] P. Tukia. Convergence groups and gromov's metric hyperbolic spaces. *New Zealand J. Math.*, 23(2):157–187, 1994.
- [29] X. Xie. Growth of relatively hyperbolic groups. *Proc. Amer. Math. Soc.*, 135(3):695–704, 2007.
- [30] A. Yaman. A topological characterization of relatively hyperbolic groups. *J. Reine Angew. Math.*, 566:41–89, 2004.

Author's addresses:

Raeyong Kim:

Department of Mathematics, The Ohio State University, 231 West 18th Avenue, Columbus, Ohio 43210-1174, United States.

`kimr@math.ohio-state.edu`